



REVIEW

Open Access

Board-invited review: Sensitivity and responses of functional groups to feed processing methods on a molecular basis

Peiqiang Yu^{1,2}

Abstract

In complex feed structures, there exist main chemical functional groups which are associated with nutrient utilization and availability and functionality. Each functional group has unique molecular structure therefore produce unique molecular vibration spectral profile. Feed processing has been used to improve nutrient utilization for many years. However, to date, there was little study on processing-induced changes of feed intrinsic structure and functional groups on a molecular basis within intact tissue. This is because limited research technique is available to study inherent structure on a molecular basis. Recently bioanalytical techniques: such as Synchrotron Infrared Microspectroscopy as well as Diffuse Reflectance Infrared Fourier Transform molecular spectroscopy have been developed. These techniques enable to detect molecular structure change within intact tissues. These techniques can prevent destruction or alteration of the intrinsic protein structures during processing for analysis. However, these techniques have not been used in animal feed and nutrition research. The objective of this review was show that with the advanced technique, sensitivity and responses of functional groups to feed processing on a molecular basis could be detected in my research team. These functional groups are highly associated with nutrient utilization in animals.

Keywords: Sensitivity and responses of functional groups, Molecular structures, Feed processing, Nutrient utilization and availability

Introduction

Functional groups in feeds

Each feed has a unique structure. The unique structure results in unique molecular vibrational spectra. Functional groups are associated within unique feed intrinsic structure. These functional groups consist of macromolecular such as lipid, protein, structural and non-structural carbohydrates which are nutrient for animal and human. Under infrared vibrational molecular spectroscopy, these functional groups produced unique bands [1]. From the IR spectrum ($4000\text{--}500\text{ cm}^{-1}$) the presence or absence of various organic functional groups is readily observed [2-4].

Molecular vibration of functional groups

Several relatively unique absorption of lipid vibration spectra ($4000\text{--}500\text{ cm}^{-1}$) usually locate at several regions in mid-infrared regions. (a) It located at ca. $2800\text{--}3000\text{ cm}^{-1}$, asymmetric and symmetric stretching vibrations of CH_3 (ca. 2956 and 2874 cm^{-1}) and CH_2 (ca. 2922 and 2852 cm^{-1}) groups of acyl chains [5-8]. (b) it locates at ca. 3015 cm^{-1} , unsaturation (CH attached to $\text{C}=\text{C}$). (c) It locate at ca. $1800\text{--}1500\text{ cm}^{-1}$ at ca. 1740 cm^{-1} stretching vibration of the carbonyl ester $\text{C}=\text{O}$ bond [2,5]. The protein produces relative unique bands which are amide I and amide II at ca. $1710\text{--}1480\text{ cm}^{-1}$. The amide I band arises predominantly from the $\text{C}=\text{O}$ stretching vibration of the peptide bond and particularly sensitive to and can be used to predict protein secondary structure [5-8]. The relative unique absorption bands from carbohydrates are complex. Normally carbohydrate bands are found at the $1180\text{--}900\text{ cm}^{-1}$ region of the infrared spectrum. There are some unique carbohydrate bands,

Correspondence: peiqiang.yu@usask.ca

¹Department of Animal and Poultry Science, College of Agriculture and Bioresources, University of Saskatchewan, 51 Campus Drive, Saskatoon S7N 5A8, Canada

²Professor and Ministry of Agriculture Strategic Research Chair, (教授,农业部战略研究主席), College of Agriculture and Bioresources, University of Saskatchewan, 51 Campus Drive, Saskatoon S7N 5A8, Canada



such as band at 1420 cm^{-1} for β -glucan [2], bands at ca. 1420 , 1370 and 1335 cm^{-1} for structural carbohydrates [2,3], band at ca. 1025 cm^{-1} for starch in endosperm of cereal grain [2,3], band at ca. 1240 cm^{-1} for hemicellulose. These functional groups are highly related to nutrient availability [9,10]

Molecular spectroscopy and spectral analysis

There are several molecular spectroscopy that could be used for molecular structure study for feeds. First one is cutting-edge bioanalytical technique: synchrotron-radiation based technique SR-IMS. This technique enables to detect molecular structure change within intact tissues within cellular and subcellular dimensions [11-14]. This technique can prevent destruction or alteration of the intrinsic protein structures during processing for analysis. Other molecular spectroscopy technique could be used for feed inherent structure study included ATR/FT/IR spectroscopy [15], DRIFT molecular spectroscopy [16]. For molecular spectral analysis, it includes univariate and multivariate analyses [4,17-19]. For univariate analysis, we could determine each functional group intensity and distribution within intact feeds tissue. For multivariate analysis, there are two methods are commonly used in my research team to discriminate and classify feed molecular structures and processing-induced structure changes, such as heating, autoclaving, bioethanol processing and gene transformation [20]. The first one is agglomerative hierarchical cluster analysis (AHCA). AHCA is initially used to try to determine the main sources of variation in SRFTIRM protein spectra and the results are displayed as dendrograms using Ward's method (Cytospec Software).

Review objective

However, these molecular-based techniques have not been used in animal feed and nutrition research. The objective of this review was to show that with the advanced technique, sensitivity and responses of functional groups to feed processing on a molecular basis could be detected in my research team.

Application I

Processing-induced changes in lipid-related function groups on a molecular basis

In one of our research [21], we studied heat-induced changes to lipid molecular structure in Vimy flaxseed: spectral intensity and molecular clustering. In this study, we characterized any changes of each functional groups mainly associated with lipid structure in flaxseed (*Linum usitatissimum*, cv. Vimy), that occurred on a molecular level during the treatment process using infrared molecular spectroscopy. We autoclaved flaxseed samples at 120°C for 20, 40 or 60 min for treatments 1, 2 and 3, respectively. The lipid CH_3 asymmetric (ca. 2959 cm^{-1}),

CH_2 asymmetric (ca. 2928 cm^{-1}), CH_3 symmetric (ca. 2871 cm^{-1}) and CH_2 symmetric (ca. 2954 cm^{-1}) functional groups, lipid carbonyl $\text{C}=\text{O}$ ester group (ca. 1745 cm^{-1}), lipid unsaturation group (CH attached to $\text{C}=\text{C}$) (ca. 3010 cm^{-1}) and their intensity ratios were determined. We also used hierarchical cluster analysis (CLA) and principal components analysis (PCA) to identify molecular structure difference. The results showed that there were linear and quadratic effects of the treatment time from 0, 20, 40 and 60 min on the ratios of the CH_2 asymmetric to CH_3 asymmetric stretching vibration intensity. But autoclaving had no significant effect on lipid carbonyl $\text{C}=\text{O}$ ester group and lipid unsaturation group (CH attached to $\text{C}=\text{C}$). The results indicated that autoclaving had different impact on different functional groups which are related to lipid and molecular spectroscopy was able to pick up heat-induced changes in lipid conformation. A future study is needed to quantify the relationship between lipid molecular structure changes and functionality/availability [21]. In another study [22], we reported sensitivity of molecular functional groups to bioethanol processing in lipid biopolymer of co-products using DRIFT molecular spectroscopy.

Application II

Processing-induced changes in protein-related function groups on a molecular basis

The sensitivity and responses of protein-related functional group to various processing has been reviewed in details and published in Molecular Nutrition and Food [20]. In this article we have reported the effects of various processing included gene-transformation, bioethanol processing, autoclaving and dry roasting on protein molecular structure changes in feeds. We also reported the responses of each functional group to the processing. The results show, again, different functional groups have different sensitivity to different processing. In Yu et al. [23], study, we used advanced synchrotron technique- SR-IMS to compare protein molecular structure of alfalfa plant tissues transformed with the maize *Lc* regulatory gene with non-transgenic alfalfa protein within cellular and subcellular dimensions and to quantify protein inherent structure profiles using Gaussian and Lorentzian methods of multi-component peak modeling [17,24]. It was found that transgenic *Lc* alfalfa leaves contain similar proteins to non-transgenic alfalfa (because amide I and II intensities were identical), but a subtle differences in protein molecular structure. Further study is needed to understand the relationship between these structural profiles and biological features such as protein nutrient availability, protein bypass and digestive behavior of livestock fed with this type of forage [23]. In the study by Doiron et al. [25], it was found that autoclaving changed protein structure α -helix to β -sheet ratio and protein sub-fractions. The

protein structure α -helix to β -sheet ratio had significantly positive correlation with total intestinally absorbed protein supply and negative correlation with degraded protein balance [25]. In the study bioethanol processing study, it was found that bioethanol processing changed protein molecular structure α -helix to β -sheet ratio and protein amide I to II ratio which highly related to nutrient values. The results indicated that the protein structure α -helix to β -sheet ratio in the new co-products of bioethanol productions are correlated to total intestinally absorbed protein supply.

Application III

Processing-induced changes in carbohydrate-related function groups on a molecular basis

Feed processing has different impact on different functional groups in feeds which are related to carbohydrate. Yu [26], reported the study on effect of bioethanol processing on carbohydrate functional groups by using DRIFT molecular spectroscopy. The following functional groups were studied, including: "A_Cell (peaks area region and baseline: ca. 1485–1188 cm^{-1}) mainly associated with hemi- and cellulosic compounds, A_1240 (peak area centered at ca.1240 cm^{-1} with region and baseline: ca. 1292–1198 cm^{-1}) mainly associated with cellulosic compounds, A_CHO (peaks region and baseline: ca. 1187–950 cm^{-1}) associate with total CHO, A_928 (peak area centered at ca. 928 cm^{-1} with region and baseline: ca. 952–910 cm^{-1}) and A_860 (peak area centered at ca. 860 cm^{-1} with region and baseline: ca. 880–827 cm^{-1}) mainly associated with non-structural CHO, H_1415 (peak height centered at ca.1415 cm^{-1} with baseline: ca. 1485–1188 cm^{-1}) and H_1370 (peak height at ca.1370 cm^{-1} with a baseline: ca. 1485–1188 cm^{-1}) mainly associated with structural CHO". The authors found that A_Cell, A_CHO, H_1415 and H_1370 had no correlation with CHO chemical and nutrient profiles in the bioethanol co-products. However, CHO molecular spectral intensity of the A_1240, A_928 and A_860 had strong correlation with rapidly degradable CHO fraction (CB1), lowly degradable CHO fraction (CB2) and an unavailable CHO fraction (CC) and could be a good indicator. In conclusion, the changes of the CHO molecular structures during the processing for bioethanol production were highly associated with carbohydrate degradable subfractions in ruminants.

Conclusions and implications

The review shows that molecular spectroscopy technique could be used to study the sensitivity and response of functional groups in complex feed systems to various feed processing on molecular basis. It has a potential to study processing impact directly on feed and nutrient utilization and availability. These functional groups are highly associated with nutrient utilization in animals.

Competing interests

The authors declare that they have no competing interests.

Acknowledgments

The chair (PY) research programs have been supported by grants from the Natural Sciences and Engineering Research Council of Canada (NSERC), SaskCanola Funding, Saskatchewan Agricultural Development Fund (ADF) and Ministry of Agriculture Strategic Research Chair Fund (Saskatchewan, Canada).

Received: 26 July 2012 Accepted: 6 December 2012

Published: 10 December 2012

References

1. Kemp W: *Organic spectroscopy*. 3rd edition. New York, USA: W.H. Freeman and Company; 1991.
2. Wetzel DL, Eilert AJ, Pietrzak LN, Miller SS, Sweat JA: **Ultraspatially resolved synchrotron infrared microspectroscopy of plant tissue in situ.** *Cell Mol Bio* 1998, **44**:145–167.
3. Wetzel DL, Srivarin P, Finney JR: **Revealing protein infrared spectral detail in a heterogeneous matrix dominated by starch.** *Vib Spectrosc* 2003, **31**:109–114.
4. Miller LM: *Infrared Microspectroscopy and Imaging*. Available: <http://www.bnl.gov/ps/nsls/newsroom/publications/otherpubs/imaging/workshopmiller.pdf>.
5. Marinkovic NS, Huang R, Bromberg P, Sullivan M, Toomey J, Miller LM, Sperber E, Moshe S, Jones KW, Chouparova E, Lappi S, Franzen S, Chance MR: **Center for Synchrotron Biosciences' U2B beamline: an international resource for biological infrared spectroscopy.** *J Synchrotron Rad* 2002, **9**:189–197.
6. Jackson M, Haris PI, Chapman D: **Conformational transitions in poly-L-lysine: an FT-IR spectroscopic study.** *Biochim Biophys Acta* 1989, **998**:75.
7. Jackson M, Mantsch HH: **The use and misuse of FTIR spectroscopy in the determination of protein structure.** *Biochem and Mol Biol* 1995, **30**:95–120.
8. Jackson M, Mantsch HH: **Biomedical Infrared Spectroscopy.** In *Pages 311–340 in Infrared Spectroscopy of Biomolecules*. Edited by Mantsch HH, Chapman D. New York: Wiley-Liss; 1996.
9. Yu P: **Protein secondary structures (α -Helix and β -Sheet) at a cellular level and protein fractions in relation to rumen degradation behaviors of protein: A novel approach.** *British J Nutri* 2005, **94**:655–665.
10. Yu P, Nuez-Ortin WG: **Relationship between protein molecular structures and metabolisable proteins in different types of dried distillers grains with soluble: a novel approach.** *Br J Nutri* 2010, **104**:1429–1437.
11. Yu P: **Application of advanced synchrotron-based Fourier transform infrared microspectroscopy (SR-FTIR) to animal nutrition and feed science: a novel approach.** *Br J Nutri* 2004, **92**:869–885.
12. Yu P: **Molecular chemistry imaging to reveal structural features of various plant feed tissues.** *J Structural Biol* 2005, **150**:81–89.
13. Yu P, Doiron K, Liu D: **Shining light on the difference in molecular structural chemical make-up and the cause of distinct biodegradation behavior between malting- and feed-type barley: A novel approach.** *J Agric Food Chem* 2008, **56**:3417–3426.
14. Liu N, Yu P: **Characterize microchemical structure of seed endosperm within a cellular dimension among six barley varieties with distinct degradation kinetics, using ultraspatially resolved synchrotron-based infrared microspectroscopy.** *J Agric Food Chem* 2010, **58**:7801–7810.
15. Samadi Y: P: **Determine heat-induced changes of protein molecular structure, protein subfraction and nutrient profiles in soybean seeds affected dry and moisture heat processing.** *J Dairy Sci* 2011, **94**:6092–6102.
16. Liu N, Yu P: **Using DRIFT molecular spectroscopy with uni- and multi-variate molecular spectral techniques to detect plant protein molecular structure difference among different genotypes of barley.** *J Agric Food Chem* 2010, **58**:6264–6269.
17. Yu P: **Multi-component peak modeling of protein secondary structures: comparison of Gaussian with Lorentzian analytical method for plant feed and seed molecular biology and chemistry research.** *Appl Spectrosc* 2005, **59**:1372–1380.
18. Marinkovic NS, Chance MR: **Synchrotron Infrared Microspectroscopy.** In *Encyclopedia of Molecular Cell Biology and Molecular Medicine, Volume 13*. 2nd edition. Edited by Meyers R: Wiley Inc; 2006:671–708.

19. Miller LM, Dumas P: **Chemical imaging of biological tissue with synchrotron infrared light.** *Biochim Biophys Acta* 2006, **1758**:846–857.
20. Yu P: **Plant-based food and feed protein structure changes induced by gene-transformation, heating and bio-ethanol processing: A novel synchrotron-based molecular structure and nutrition research program.** *Mol Nutri Food Res* 2010, **54**:1535–1545.
21. Yu P, Damiran D: **Heat-induced changes to lipid molecular structure in Vimy flaxseed: Spectral intensity and molecular clustering.** *Spectrochim Acta A Mol Biomol Spectrosc (Elsevier)* 2011, **79**:51–59.
22. Yu P: **Study the Sensitivity of molecular functional groups to bioethanol processing in lipid biopolymer of co-products using DRIFT molecular spectroscopy.** *Spectrochim Acta A Mol Biomol Spectrosc (Elsevier)* 2011, **82**:1–7.
23. Yu P, Jonker A, Gruber M: **Molecular basis of protein structure and nutritive value in proanthocyanidin-enhanced Lc-transgenic alfalfa using synchrotron-radiation FTIR microspectroscopy.** *Spectrochim Acta A Mol Biomol Spectrosc (Elsevier)* 2009, **73**:846–853.
24. Yu P: **Synchrotron IR Microspectroscopy for Protein Structure Analysis: Potential and Questions: A Review.** *Spectroscopy* 2006, **20**:229–251. Invited Review Article.
25. Doiron KJ, Yu P, McKinnon JJ, Christensen DA: **Heat-induced protein structures and protein subfractions in relation to protein degradation kinetics and intestinal availability in dairy cattle.** *J Dairy Sci* 2009, **92**:3319–3330.
26. Yu P: **Relationship of carbohydrate molecular spectroscopic features to carbohydrate nutrient profiles in new co-products from bioethanol production.** *J Dairy Sci* 2012, **95**:2091–2096.

doi:10.1186/2049-1891-3-40

Cite this article as: Yu: Board-invited review: Sensitivity and responses of functional groups to feed processing methods on a molecular basis. *Journal of Animal Science and Biotechnology* 2012 **3**:40.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

